

Strong curvature singularities in quasi-spherical asymptotically de Sitter dust collapse

Sérgio M C V Gonçalves

Theoretical Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA

Received 5 June 2001, in final form 21 August 2001

Published 22 October 2001

Online at stacks.iop.org/CQG/18/4517

Abstract

We study the occurrence, visibility, and curvature strength of singularities in dust-containing Szekeres spacetimes (which possess no Killing vectors) with a positive cosmological constant. We find that such singularities can be locally naked, Tipler strong and develop from a non-zero-measure set of regular initial data. When examined along timelike geodesics, the singularity's curvature strength is found to be independent of the initial data.

PACS numbers: 0420D, 0420J, 0470B

1. Introduction

It has recently been shown that a central, locally naked, Tipler strong singularity develops in the gravitational collapse of an inhomogeneous spherical dust with a positive cosmological constant [1, 2]. Those results are a generalization to asymptotically de Sitter spacetimes of the well known asymptotically flat spherical dust (Lemaître–Tolman–Bondi (LTB)) collapse models. The exact solvability of such models enables a detailed analysis of gravitational collapse, and formation and structure of singularities, which shows that, from regular initial data, a Tipler strong curvature singularity always develops, which can be locally or globally naked, depending on the initial data [3]. Although almost a paradigm for gravitational collapse, LTB models rely on the simplifying assumptions of irrotational dust and spherical symmetry. Within spherical symmetry, various generalizations of the matter model have been considered [4–8], with similar results: a central curvature singularity forms, and its visibility (local versus global) depends on the initial data.

Comparatively, much less is known about singularity formation and structure in non-spherical collapse. Thorne's seminal analysis of cylindrical collapse [9] led him to formulate the *hoop conjecture*, which essentially states that horizons form if and only when the gravitational mass of the system is confined to a maximum 'radius' in every direction [10]. Subsequent numerical analyses of prolate and oblate collapse by Shapiro and Teukolsky [11, 12], and of gravitational radiation emission in aspherical collapse by Nakamura *et al* [13], were unable to refute Thorne's conjecture. A perturbative analysis of LTB collapse by Hirada

et al [14–16], showed that linear non-spherical matter, metric and matter coupled to metric perturbations remain bounded in the limit of approach to the Cauchy horizon. The very similar structure of the central singularity in LTB and Tolman–Bondi–de Sitter (TBdS) models [1, 17], strongly suggests that analogous marginal stability properties also hold for the latter case. The quasi-spherical dust collapse models, given by the Szekeres metric [18], were analysed by Szekeres himself [19], Joshi and Królak [20] and by Deshingkar *et al* [21]. As in the spherical LTB case, a ‘central’ singularity forms, which is Tipler strong, and can be locally or globally naked, depending on the initial data.

Recent measurements of type Ia supernovae [22, 23] and peculiar motion of low-redshift galaxies [24], appear to suggest the existence of a positive cosmological constant. The possibility of a non-zero asymptotic (constant) curvature constitutes an obvious motivation for the study of gravitational collapse with a cosmological constant [1, 2, 25]. In this paper, we address one aspect of this problem by studying the formation, visibility and curvature strength of singularities in quasi-spherical inhomogeneous spacetimes with a positive cosmological constant, thereby generalizing the results of [1] to the quasi-spherical case, and those of [19–21] to the asymptotically de Sitter case.

The general metric for Szekeres spacetimes with a positive cosmological constant was obtained in closed form by Barrow and Stein-Schabes [26], in the context of the ‘cosmic no-hair theorem’, but this solution also provides a valuable test-bed metric for the analysis of singularity formation and structure in non-spherical gravitational collapse. One must remark, however, that such a departure from spherical symmetry, far from being arbitrary, is very well defined—whilst the four-dimensional spacetime metric does not admit any Killing vectors, it does possess an invariant family of spherical 2-surfaces, hence the name *quasi-spherical*. Accordingly, the results presented here provide a useful, but somewhat limited, insight into non-perturbative departures from spherical symmetry. Note also that, Szekeres spacetimes can be matched to a Schwarzschild (or Schwarzschild–de Sitter, if $\Lambda > 0$) spacetime, and thus cannot contain gravitational waves [27].

As we shall see below, as in the Szekeres and TBdS cases, for quasi-spherical asymptotically de Sitter dust collapse, a central singularity develops from regular initial data, and is at least locally naked and Tipler strong. Our results suggest that neither lack of asymptotic flatness, nor ‘mild’ deviations (in a sense to be precisely defined below) from spherical symmetry, play an important role in the formation and nature of singularities in gravitational collapse.

The paper is organized as follows. Szekeres spacetimes with a cosmological constant are outlined in section 2. Section 3 discusses the existence and visibility of the singularity, along null and timelike directions. The curvature strength of the singularity is analysed in section 4. Section 5 concludes with a discussion and summary.

Geometrized units, in which $G = c = 1$, are used throughout.

2. Szekeres spacetimes with a cosmological constant

In this section we present the relevant equations for Szekeres metrics [18, 19] with a positive cosmological constant [26]. The stress–energy tensor is

$$T_{ab} = (p + \rho)u_a u_b - \left(p - \frac{\Lambda}{8\pi}\right)g_{ab}, \quad u_a = \delta_a^t, \quad (1)$$

where u_a , p and ρ are the 4-velocity, pressure and fluid density, respectively. The metric can be written in normal Gaussian coordinates:

$$ds^2 = -dt^2 + e^{2\alpha} dr^2 + R^2 e^{2\nu} (dx^2 + dy^2), \quad (2)$$




Figure 1. Conformal diagram for asymptotically de Sitter quasi-spherical dust collapse, for the 2-surface $\Sigma : \{x, y = \text{constant}\}$. The $r = 0$ shell is the first one to become singular. The outer shell, which defines the boundary of Σ , first collapses through the past cosmological horizon (H^-), and then, after a finite amount of time $t = t_c$, through the black hole event horizon (EH). The example depicted here contains a locally naked singularity, whose null portion is visible to local observers inside the EH. Since the EH coincides with the Cauchy horizon, the singularity cannot be globally naked.

where $\alpha = \alpha(t, r, x, y)$, $R = R(t, r)$, $v = v(r, x, y)$ and

$$e^{\alpha(t, r, x, y)} = [R'(t, r) + R(t, r)v'(r, x, y)][1 + k(r)]^{-1/2}, \quad (3)$$

$$e^{-v(r, x, y)} = c_1(x^2 + y^2) + 2(c_2x + c_3y) + c_4, \quad (4)$$

$$k(r) = 4(c_1c_4 - c_2^2 - c_3^2) - 1, \quad (5)$$

where $' \equiv \partial_r$, $\dot{} \equiv \partial_t$ and the real-valued functions $c_i = c_i(r)$, $i = 1 \dots 4$, are to be specified within each Szekeres class. The Szekeres metrics (with or without a cosmological constant) can be divided into two classes, depending on whether $(Re^v)'$ vanishes or not [18, 26]. The class defined by $(Re^v)' = 0$ contains shell-crossing singularities [28, 29], which are physically ‘mild’—they are gravitationally weak [30] and geodesically complete [31]—and hence will not be considered here. We shall focus on the class for which $(Re^v)' \neq 0$, which admits shell-focusing curvature singularities, but no shell-crossings. An example of an asymptotically de Sitter quasispherical spacetime containing such a shell-focusing singularity is shown in figure 1.

The function $R(t, r)$ obeys the evolution equation

$$\dot{R}^2 = \frac{2M(r)}{R} - k(r) + \frac{1}{3}\Lambda R^2, \quad (6)$$

where $M(r)$ is a real-valued free function. Introducing an auxiliary variable η , defined by

$$\sqrt{R} d\eta = dt. \quad (7)$$

Equation (6) reduces to

$$\frac{d\eta}{d\theta} = \pm \sqrt{\frac{6}{\Lambda p}} (\gamma \pm \cosh \theta)^{-1/2}, \quad (8)$$

which has solutions of the form

$$\eta = \eta_0 \pm 2 \sqrt{\frac{6}{\Lambda p}} (\gamma + 1)^{-1/2} F(\psi, a), \quad (9)$$

where $\eta_0 = \eta(\theta = 0)$ and $F(\psi, a)$ is an elliptic function of the third kind [32], with $\psi \equiv \sin^{-1} \Psi$, where

$$\Psi \equiv \tanh(\theta/2) \quad ('+' \text{ in } (8)), \quad (10)$$

$$\Psi \equiv \sqrt{(\gamma - \cosh \theta)/(\gamma - 1)} \quad ('-' \text{ in } (8)), \quad (11)$$

and

$$4 \left(R^2 + R\chi - \frac{3M}{\Lambda\chi} \right) = \left(\chi^2 - \frac{12M}{\Lambda\chi} \right) \sinh^2 \theta, \quad (12)$$

where

$$\gamma \equiv -3 \frac{\chi}{p}, \quad (13)$$

$$\chi = \begin{cases} -|4k/\Lambda|^{1/2} \cosh \Theta, & k \geq 0, \\ -|-4k/\Lambda|^{1/2} \sinh \Theta, & k \leq 0, \end{cases} \quad (14)$$

$$\Theta \equiv \frac{1}{3} \cosh^{-1} \left(\frac{3M}{2} \sqrt{\left| \frac{-\Lambda}{k^3} \right|} \right), \quad (15)$$

$$p \equiv \sqrt{\chi^2 + \frac{12M}{\Lambda\chi}}, \quad (16)$$

$$a \equiv \sqrt{(\gamma - 1)/(\gamma + 1)}. \quad (17)$$

For the particular case of dust ($p = 0$), with gravitationally unbound matter distributions ($k = 0$), the solution reduces to the simple algebraic form:

$$R(t, r) = \left(\frac{3M}{\Lambda} \right)^{1/3} \sinh^{2/3} \left\{ \sqrt{\frac{3\Lambda}{4}} [t_c(r) - t] \right\}, \quad (18)$$

where t_c is a real-valued arbitrary function, fixed by the initial data via

$$t_c(r) = \frac{2}{\sqrt{3\Lambda}} \sinh^{-1} \left(\sqrt{\frac{\Lambda r^3}{3M}} \right), \quad (19)$$

where the scaling $R(0, r) = r$ was adopted. The relevant derivatives of $R(t, r)$ are

$$R'(t, r) = R \left[\frac{M'}{3M} + \sqrt{\frac{\Lambda}{3}} t'_c \coth(t_c - t) \right], \quad (20)$$

$$\dot{R}(t, r) = -\sqrt{\frac{\Lambda}{3}} R \coth(t_c - t), \quad (21)$$

where the minus sign corresponds to implosion.

A complete solution is given by equation (3), where R and ν are given by equations (4), (5) and (18), together with the energy density

$$\rho = (M' - 3M\nu')e^{-\alpha-2\beta} = \frac{M' - 3M\nu'}{R^2(R' + R\nu')}e^{-2\nu}. \quad (22)$$

Imposing the weak energy condition [33], $T_{ab}u^a u^b \geq 0 \Rightarrow \rho \geq 0$, leads to the constraint

$$\nu' \leq \frac{1}{3} \frac{M'}{M}. \quad (23)$$

2.1. Regularity conditions

We assume the following regularity conditions for the spacetime.

- (a) There are no shell-crossing singularities: $(Re^\nu)' > 0$, which also guarantees that the proper area radius of the shells is a monotonically increasing function of the comoving spatial coordinate r ,

$$R' + R\nu' > 0. \quad (24)$$

- (b) Regularity of the metric at $r = 0$ requires that all the $c_i(r)$ be at least C^1 therein, with finite first derivatives, such that

$$\lim_{r \rightarrow 0^+} c'_i(r) < \infty \Rightarrow \lim_{r \rightarrow 0^+} \nu' < \infty. \quad (25)$$

- (c) In order for the metric to be locally Euclidean at $r = 0$, we must have

$$k(0) = 0. \quad (26)$$

- (d) From a physical point of view, it is reasonable to require that the initial velocity profile, $\dot{R}(0, r)$, and the energy density, $\rho(0, r, x, y)$, be everywhere bounded. With the scaling $R(0, r) = r$, it follows from equation (6) that $M \gtrsim O(r)$ near $r = 0$ and equation (22) implies $M' \sim O(r^2)$; near $r = 0$ we must therefore have

$$\frac{M'}{M} \sim O(r^{-1}). \quad (27)$$

3. Existence and visibility

Each 2-surface $\Sigma : \{t, r = \text{constant}\}$ is a 2-sphere with proper area radius $R(t, r)e^{\nu(r, x, y)}$ and coordinate centre $(c_2/c_1, c_3/c_1)$ on Σ . Since the centre is r -dependent, the shells are not in general concentric, which accounts for the absence of spherical (or otherwise) symmetry. Note also that, unlike in the spherically symmetric case, the energy density is not constant over each 2-sphere. If three of the c_i s are constant, there is a single c_i free function—which fixes $k(r)$, via equation (5)—and the model reduces to the TBdS case (where the $k = 0$ case corresponds to the choice $c_1 = c_4 = \frac{1}{2}$ and $c_2 = c_3 = 0$).

From equation (22), one sees that the energy density diverges at $R = 0$, thereby signalling a curvature singularity [19]. As in the spherical case, $r > 0$ events for which $R(r, t) = 0$, are spacelike and hence cannot be naked [19, 20, 34]. Of potential interest is the ‘central’ $r = 0$ singularity.

In order for this singularity to be at least locally naked, there must exist at least one outgoing non-spacelike geodesic with past endpoint at the singularity. We note that the existence of a solution of the, for example, outgoing radial null geodesics equation, does

not *per se* guarantee the local visibility of the singularity. If such geodesics are emitted after the apparent horizon (AH) forms, they will be contained in the causal past of a trapped region (and hence unavoidably trapped), and thus any initially diverging geodesic congruence will immediately start reconverging upon emission. Therefore, although there may be outgoing non-spacelike geodesics with a past endpoint at the singularity, if they are emitted after the AH forms, they are only defined at the point of emission, and hence we shall not consider them in the operational definition of local nakedness adopted in this paper. We shall show that there are outgoing non-spacelike geodesics that form before or at the time at which the AH does, by analysing the geodesic behaviour along null and timelike directions.

3.1. Null geodesics

The equation for outgoing radial null geodesics (ORNGs) is

$$\left(\frac{dt}{dr}\right)_{\text{out}} = R' + Rv'. \quad (28)$$

If it admits a regular solution, $t(r)$, in an open neighbourhood containing $r = 0$, the singularity is visible—at least locally—provided $t(0) \leq t_{\text{AH}}(0)$, where $t_{\text{AH}}(r)$ is the AH curve on the $\{t, r\}$ plane. Let us then assume that such a solution exists, and is given to leading order by

$$t(r) = t_0 + ar^\sigma, \quad (29)$$

where $a \in \mathbb{R}^+$ and $\sigma \in \mathbb{N}^+$. Expanding the central energy density near $r = 0$ as

$$\rho_c(r) = \sum_{i=0}^{+\infty} \rho_i r^i, \quad (30)$$

we have, from equations (22), (23), (27), (19) and (6),

$$M(r) = M_0 r^3 + M_n r^{n+3} + O(r^{n+4}), \quad (31)$$

$$t_c(r) = t_0 + t_n r^n + O(r^{n+1}), \quad (32)$$

$$R(t, r) = \left(\frac{9}{4}\right)^{1/3} (M_0^{1/3} r + m_n r^{n+1}) (t_0 + t_n r^n - t)^{2/3} + O(r^{n+2+\sigma}), \quad (33)$$

where t_n and m_n are real coefficients linear in $M_n = (4\pi/n)\rho_n$, with $n > 0$; $\rho_n \equiv (\partial^n \rho_c / \partial r^n)_{r=0}$ is the first non-vanishing derivative of the central energy density distribution and

$$t_0 = \sinh^{-1} \left(\sqrt{\frac{\Lambda}{6M_0}} \right). \quad (34)$$

Since the geodesic must lie on the spacetime, from equations (29) and (32) it follows that $\sigma \geq n$. If $\sigma = n$, we have the additional constraint, $a < t_n$.

The existence of a self-consistent solution of equation (28) is dependent on the r -differentiability of R and v at $r = 0$. From equation (4), we have

$$v'(r, x, y) = -\frac{c'_1(x^2 + y^2) + 2(c'_2x + c'_3y) + c'_4}{c_1(x^2 + y^2) + 2(c_2x + c_3y) + c_4}. \quad (35)$$

Since all the $c_i(r)$ are assumed to be at least C^1 , to leading order their MacLaurin expansion is

$$c_i(r) = c_i^0 + c_i^{n_i} r^{n_i}, \quad (36)$$

where $c_i^{n_i} \equiv (\partial^{n_i} c_i / \partial r^{n_i})_{r=0}$ is the first non-vanishing derivative of $c_i(r)$ and $n_i \geq 1$. Without loss of generality, we can take $n_1 = \min\{n_i, i = 1, \dots, 4\}$, such that near $r = 0$ we have

$$v' = \frac{n_1 r^{n_1-1}}{r^{n_1} + \zeta}, \quad (37)$$

$$\zeta(x, y) \equiv \frac{c_1^0}{c_1^{n_1}} + (2c_2^0 x + 2c_3^0 y + c_4^0) c_1^{-n_1} (x^2 + y^2)^{-1}. \quad (38)$$

If $\zeta = 0$ (which is a set of measure zero in the initial data, since it requires $c_i^0 = 0, i = 1, \dots, 4$), then $v' = n_1 r^{-1}$ and the metric is just C^0 at $r = 0$. In general, $\zeta \neq 0$, which renders the metric C^1 at $r = 0$, as desired. In this case, from equations (28), (29) and (33), we obtain, near $r = 0$

$$a\sigma r^{\sigma-1} = \left(\frac{2n}{3} + 1\right) A r^{\frac{2}{3}n} + \frac{A}{\zeta} n_1 R^{\frac{2}{3}n+n_1} + O(r^{\sigma+1-\frac{1}{3}n}), \quad (39)$$

$$A \equiv \left(\frac{9M_0}{4}\right)^{1/3} t_n^{1/3}. \quad (40)$$

Since $n_1 \geq 1$, the second term on the right-hand side is to be neglected to leading order in r , and a self-consistent solution exists provided

$$a = \left(\frac{9M_0}{4}\right)^{1/3} t_n^{1/3} \left(\frac{2n}{3} + 1\right), \quad (41)$$

$$\sigma = 1 + \frac{2n}{3}. \quad (42)$$

The condition $\sigma > n$ now reads $n < 3$. For $n = 1, 2$ (i.e. for $\rho_1 \neq 0$, or $\rho_1 = 0$ and $\rho_2 \neq 0$) there is a self-consistent solution to the ORNG equation in the limit $t \rightarrow t_0, r \rightarrow 0$, and thus there is at least one ORNG starting from the singularity.

Let us now examine the case $n = \sigma = 3$. Proceeding as before, we obtain, to leading order,

$$3ar^2 = 3 \left(\frac{9M_0}{4}\right)^{1/3} (t_3 - a)^{2/3} r^2, \quad (43)$$

which is identically satisfied provided

$$a^3 - \mu^3 a^2 + 2\mu^2 t_3 a = 0, \quad (44)$$

where $\mu \equiv (9M_0/4)^{1/3}$. This equation has two non-zero distinct roots (other than the $a = 0$ trivial root), given by $a = (\mu^2/2) \pm \sqrt{\mu^4 - 8t_3}$, if $t_3 < \frac{1}{8} \left(\frac{9}{4}M_0\right)^{4/3}$, which imposes a constraint on ρ_3 , for a given ρ_0 . In addition, one must also require $a < t_3$, which leads to

$$\frac{\mu^2}{2} - 4 + \sqrt{\mu^4 + 16 - 4\mu^2} < t_3 < \frac{1}{8}\mu^4. \quad (45)$$

Hence, as long as one restricts ourselves to initial data that satisfies the above condition, there exists an ORNG with a past endpoint at the singularity. We note, however, that the $n = 3$ case is less generic than the $n < 3$ cases, as it requires that $\rho_1 = \rho_2 = 0$ and ρ_3 obey condition (45).

Summarizing, for a given initial density profile, the ORNG equation admits a regular solution at $r = 0$, independently of the details of $v(r, x, y)$ (provided v obeys the regularity condition (25)).

3.2. Timelike geodesics

Let us consider radial timelike geodesics (RTGs) described by a tangent vector field $K^a = dx^a/d\tau$, where τ is an affine parameter along the geodesic. A sufficient set of equations for K^a is

$$K^t = \pm \sqrt{1 + e^{2\alpha}(K^r)^2}, \quad (46)$$

$$(K^r)' + \frac{K^r}{K^t}(K^r)'R' + K^r \frac{\dot{R}' + \dot{R}v'}{R' + Rv'} + \frac{(K^r)^2}{K^t} \frac{R'' + R'v' + Rv''}{R' + Rv'} = 0, \quad (47)$$

where the first equation is the unit-norm condition, and the second one follows from the geodesic equation. By inspection, one sees that the above set admits the trivial solution

$$K^t = \pm 1, \quad (48)$$

$$K^r = 0, \quad (49)$$

which leads to

$$t = t_0 \pm (\tau - \tau_0), \quad (50)$$

$$r = r_0 = \text{constant}, \quad (51)$$

where τ_0 is the proper time at which RTGs depart (arrive at) the central singularity, and the plus or minus sign refers to outgoing or ingoing RTGs, respectively. The outgoing RTG departing from the singularity is given by $r = 0$, and $t = t_0 + \tau - \tau_0$, and thus does not belong in the spacetime. The ingoing RTG is given by $r = 0$, $t = t_0 - \tau + \tau_0$, where $t_0 = t_c(0) = 0$ is the time at which the RTG arrives at the singularity.

Since equation (47) is a mixed first-order linear PDE for $K^r(t, r)$, its solution need not be unique [35]. Indeed, one can explicitly construct other families of solutions, as follows. Near the singularity, we can write, to leading order

$$t_{\text{RTG}}(r) = t_0 + br^p, \quad (52)$$

$$R(t_{\text{RTG}}(r), r) = a_0 r^{\frac{2}{3}n+1} + O(r^{p+2-\frac{1}{3}n}), \quad (53)$$

$$R'(t_{\text{RTG}}(r), r) = a_0 \left(\frac{2n}{3} + 1 \right) r^{2n/3} + O(r^{p+1-\frac{1}{3}n}), \quad (54)$$

where $a_0 \in \mathbb{R}^+$, $b \in \mathbb{R}$ and $p \geq n$ (such that the geodesics thus constructed belong in the spacetime).

Let us now assume that $K^r(t, r) \propto (t - t_0)^\alpha r^\beta$, where $\alpha, \beta \in \mathbb{R}$. From equation (52), along the RTG, we have then

$$K^r(t_{\text{RTG}}(r), r) = kr^{\alpha p + \beta}, \quad (55)$$

where $k \in \mathbb{R}$ is a constant, yet to be determined. With these ansätze, the RTG equation reads

$$\frac{dt_{\text{RTG}}}{dr} = bpr^{p-1} = \frac{K^t}{K^r} = \sqrt{(K^r)^{-2} + (R' + Rv')^2}. \quad (56)$$

Now, from equation (37), we have $v' = n_1 \xi^{-1} r^{n_1-1}$, where $n_1 \geq 1$ is fixed by the initial data, and characterizes the differentiability of the metric at the origin. From equation (56), it follows that

$$b^2 p^2 r^{2(p-1)} = k^{-2} r^{-2(\alpha p + \beta)} + a_0^2 q^2 r^{2l} + a_0^2 \frac{n_1^2}{\xi^2} r^{2(q+n_1-1)} + 2a_0^2 q \frac{n_1}{\xi} r^{q+l+n_1-1}, \quad (57)$$

where $q = 1 + 2n/3$ and $l = 2n/3$. Now, since $n_1 \geq 1$, a straightforward calculation shows that if $n < 6$, the last two terms are of higher order than the remaining ones. We shall assume

this for now, and conclude below that a self-consistent solution always satisfies this condition. We then look for solutions which are homogeneous in the leading order of r . This leads to a coupled algebraic system for the parameters p, q, α, β :

$$p - 1 = -\alpha p - \beta, \quad (58)$$

$$\alpha p + \beta = -q + 1, \quad (59)$$

which is solved by

$$p = 1 + \frac{2n}{3} = \frac{1 - \beta}{1 + \alpha}. \quad (60)$$

The constraint $p \geq n$ now reads $n \leq 3$, which is consistent with the earlier assumption $n < 6$. For parameter values satisfying equation (60), equation (56) becomes

$$C(a_0, b, p, k)r^{4n/3} = 0, \quad (61)$$

whose solution is given by the algebraic constraint

$$C(a_0, b, p, k) \equiv b^2 p^2 - k^{-2} - a_0^2 \frac{4n^2}{9} = 0. \quad (62)$$

Now, since K^r is obtained from the derivatives of $R(t, r)$, the parameter k is not independent of a_0 ; therefore, for given initial data, the set of parameters $\{a_0, p, k\}$ is uniquely determined. From equation (62), it then follows that b is entirely determined from the initial data, modulo its sign, i.e. for a given set of initial data, both ingoing ($b < 0$) and outgoing ($b > 0$) radial timelike geodesics exist.

3.3. Local visibility

Thus far we have shown the existence of outgoing null and timelike geodesics with past endpoint at the singularity. In order for these geodesics to be visible to local non-spacelike observers, they must not be contained in the domain of dependence of any trapped surfaces. Clearly, a necessary and sufficient condition for the singularity to be locally naked is that future-directed geodesics departing from the singularity do so before or at the time at which the AH forms; otherwise they will be unavoidably trapped.

The AH is the outer boundary of a trapped surface—a compact spacelike 2-surface whose outgoing and ingoing null geodesic congruences have vanishing expansion. Let us then consider the 2-surface $\Sigma : \{t, r = \text{constant}\}$, and a congruence of radial null geodesics, with tangent vector field $\xi^a = dx^a/d\lambda$ (where λ is an affine parameter along the geodesic), orthogonal to it. The expansion of such a congruence is given by the scalar

$$\Theta \equiv \nabla_a \xi^a. \quad (63)$$

Orthogonality to Σ allows the choice

$$\xi^x = \xi^y = 0, \quad (64)$$

and the null-norm condition gives

$$(\xi^t)^2 = (R' + R\nu')^2 (\xi^r)^2. \quad (65)$$

One can choose the affine parameter such that

$$\xi^t = (R' + R\nu')^2, \quad (66)$$

$$\xi^r = \epsilon = \pm 1, \quad (67)$$

where the plus or minus sign corresponds to outgoing or ingoing geodesics, respectively. From equations (2)–(5) and (66), (67), we then have

$$\Theta = \frac{2}{R}(R' + Rv')(\dot{R} + \epsilon). \quad (68)$$

Now, from equation (24) the first term is always positive-definite, and thus $\text{sign } \Theta = \text{sign } (\dot{R} + \epsilon)$. Since we are interested in implosion situations, we must also have $\dot{R} \leq 0$ for all times. It then follows that for ingoing null geodesics ($\epsilon = -1$), we have $\Theta < 0$, i.e. their expansion is always negative (convergence). For outgoing geodesics ($\epsilon = +1$) one can have initial divergence, if $\dot{R}(0, r) > -1$. However, since¹

$$\dot{R}(0, r) = -\frac{M(r)}{r^2} + \frac{\Lambda}{3}r < 0,$$

there is a finite time $t_M(r) > 0$ such that $\dot{R}(t_M, r) < -1$. Therefore, there is a time $t_{AH} \in (0, t_M)$, such that

$$\dot{R}(t_{AH}, r) = -1. \quad (69)$$

It then follows from equation (6) that the AH is given by the curve $t_{AH}(r)$, where $R(t_{AH}, r)$ is a solution of

$$\frac{\Lambda}{3}R^3 - R + 2M = 0. \quad (70)$$

This equation has three distinct real roots if $3M\sqrt{\Lambda} < 1$, two of which are positive and given by

$$R_{\pm} = \frac{2}{\sqrt{\Lambda}} \sin \left[\frac{1}{3} \sin^{-1}(3M\sqrt{\Lambda}) + \frac{2\pi}{3} \delta(1 \pm 1) \right], \quad (71)$$

where $\delta(x)$ is the Dirac delta-function, and $R_- > R_+ > 0$, corresponding to the choice $0 \leq \sin^{-1} \omega \leq \pi/2$, $0 \leq \omega \leq 1$. The third root, $R_3 = -R_- - R_+$, is negative and hence unphysical. R_- is a generalized cosmological horizon ($R_- = \sqrt{3/\Lambda}$, when $M = 0$) and R_+ the black hole apparent horizon ($R_+ = 2M$ when $\Lambda = 0$; the apparent and event horizons coincide in the static case). For $3M\sqrt{\Lambda} = 1$, the two horizons coincide. If $3M\sqrt{\Lambda} > 1$, there is one negative real root and two complex (conjugate) roots, all of which are unphysical: the spacetime does not admit any horizons in this case. From the ‘+’ solution in equation (71), together with equations (18) and (19), we obtain

$$t_{AH}(r) = t_c(r) - \varsigma(r), \quad (72)$$

where

$$\varsigma(r) \equiv \frac{2}{\sqrt{3\Lambda}} \sinh^{-1} \left\{ \left(\frac{8}{3M\sqrt{\Lambda}} \right)^{1/3} \left[\sin \left(\frac{1}{3} \sin^{-1}(3M\sqrt{\Lambda}) \right) \right]^{3/2} \right\}. \quad (73)$$

At the origin, $\varsigma(0) = 0$ and thus $t_{AH}(0) = t_c(0)$. Since the AH and the singularity curve form at the same time at $r = 0$, the visibility of the singularity is determined by the relative slopes of the curves $t_{AH}(r)$ and $t_{ORG}(r)$ on the $\{t, r\}$ plane (where ‘ORG’ denotes non-spacelike outgoing radial geodesics). The singularity is (at least) locally naked iff

$$\lim_{r \rightarrow 0} \left(\frac{dt_{AH}}{dr} \right) / \left(\frac{dt_{ORG}}{dr} \right) > 1. \quad (74)$$

¹ Near $r = 0$ this condition becomes $\rho_c(0) \gtrsim \Lambda$, which is always satisfied for physically reasonable values of the central density and cosmological constant (in geometrized units).

For ORNGs, given by $n < 3$, from equations (29) and (42), we obtain

$$\lim_{r \rightarrow 0} \left(\frac{dt_{\text{AH}}}{dr} \right) / \left(\frac{dt_{\text{ORNG}}}{dr} \right) = \lim_{r \rightarrow 0} \frac{nt_n}{a\sigma} r^{-1+n/3} = +\infty, \quad (75)$$

and hence the singularity is locally naked. This is in agreement with the analysis of [2], which shows that the value of Λ does not alter the visibility of the singularity for $n < 3$. (For the highly non-generic case $n = 3$, part (but *not* all) of the singularity spectrum may be covered for sufficiently large (but finite) Λ ; the method discussed in the present paper is not appropriate for such an analysis, and we thus refer to reader to [2] for further details).

For ORTGs, given by equation (56), the local visibility condition reads

$$\lim_{r \rightarrow 0} \frac{nt_n}{bp} r^{n-p} > 1. \quad (76)$$

Clearly, we must have $p > \sigma > n$, which yields $\lim_{r \rightarrow 0} (nt_n/bp)r^{n-p} = +\infty$. Hence, ORTGs are locally visible.




Figure 2. Locally naked singularity in quasi-spherical asymptotically de Sitter dust collapse. The broken curve is the apparent horizon (AH), which joins the event horizon (EH) when the boundary of the 2-surface Σ crosses the latter, at event P . Also shown is the worldline of a timelike observer O , who penetrates the EH and terminates at the spacelike portion of the singularity. PNC denotes the past null cone of O , which intersects the EH and the null portion of the singularity: clearly, there are non-spacelike geodesics emitted from the singularity, before the AH forms, which are contained in the PNC of O ; the singularity is therefore locally naked.

4. Curvature strength

Let $\gamma(\tau)$ be a RTG, with tangent vector $K^a = dx^a/d\tau$, that terminates at the singularity at $\tau = \tau_0$. Such a singularity is said to be Tipler strong along $\gamma(\tau)$ if the volume 3-form $V(\tau)$ vanishes in the limit $\tau \rightarrow \tau_0$ [36]. If the scalar $\Psi \equiv R_{ab}K^aK^b$ obeys the *strong limiting focusing condition*,

$$\lim_{\tau \rightarrow \tau_0} (\tau - \tau_0)^2 \Psi > 0, \quad (77)$$

then the singularity is gravitationally strong in the sense of Tipler [37]. This sufficient condition guarantees that any 3-form defined along $\gamma(\tau)$ vanishes in the limit $\tau \rightarrow \tau_0$, due to unbounded curvature growth.

From equations (2)–(5) and (48), (49), we have

$$\Psi = (1 + \xi)^{-1} \left(\frac{M'}{R^2 R'} + \frac{4M}{R^3} \xi \right) - \Lambda, \quad (78)$$

$$\xi \equiv \frac{R}{R'} v'. \quad (79)$$

Now, from equation (20) it follows that $\lim_{r \rightarrow 0^+} (R/R') = 0$, and from equation (23) we have $\lim_{r \rightarrow 0^+} v' < \infty$. Therefore, $\lim_{r \rightarrow 0^+} \xi = 0$, which implies

$$\lim_{r \rightarrow 0^+} \Psi = \lim_{r \rightarrow 0^+} \frac{M'}{R^2 R'} - \Lambda. \quad (80)$$

Using equations (31)–(33), together with equation (50), we obtain

$$\lim_{r \rightarrow 0^+} \Psi = \frac{2}{3} \eta^{-4/3} \left(\eta^{2/3} + \frac{2}{3} n t_n r^n \eta^{-1/3} \right)^{-1}, \quad (81)$$

$$\eta \equiv t_n r^n \mp (\tau - \tau_0). \quad (82)$$

Hence, at $r = 0$ we have

$$\lim_{\tau \rightarrow \tau_0} (\tau - \tau_0)^2 \Psi = \frac{2}{3}, \quad (83)$$

and the singularity is therefore Tipler strong. That is, along the timelike geodesics given by equations (48) and (49), the singularity is Tipler strong, independently of the details of the initial density distribution. We note that, since all the sufficient criteria for Tipler strong singularities are sufficient for *deformationally strong* ones [38], the central curvature singularity in TBdS collapse is also deformationally strong.

5. Concluding remarks

We found that there are initial data, for which null and timelike families of outgoing radial geodesics with past endpoint at the singularity exist, which are emitted along the apparent horizon curve; the singularity is therefore at least locally naked. When examined along timelike geodesics terminating at the singularity, the latter was found to be Tipler strong, regardless of the initial density distribution. This constitutes a rather robust result, in that it holds true independently of the initial data (the initial density profile $\rho(0, r, x, y)$, for any given velocity profile $\dot{R}(0, r) < 0$), thereby implying stability against perturbations of the latter.

It was shown in [21] that there is no directional dependence of the local visibility in asymptotically flat Szekeres dust collapse. We obtained the same result for asymptotically de Sitter quasi-spherical dust collapse, as expected. We also showed the absence of effects of Λ on the visibility of singularities for $n < 3$, already observed in spherical Tolman–Bondi–de Sitter collapse. The fact that our results are qualitatively equivalent to those for spherical inhomogeneous collapse (with or without a cosmological constant), suggests that neither the lack of asymptotic flatness, nor mild (in the well defined geometric sense of Szekeres metrics) departures from spherical symmetry change the standard picture of singularity formation and structure in gravitational collapse.

Regarding the generality of the present results insofar as the matter content is concerned, two points are worth mentioning. Firstly, we considered the special $k(r) = 0$ case (cf equation (5)), which corresponds to gravitationally unbound matter configurations. This simplifying assumption ($k = 0$) was found not to qualitatively change the corresponding results for spherical LTB collapse [39] (note, however, that in the asymptotically flat case, $k = 0$ corresponds to marginally bound matter configurations). Since the structure of the central singularity in the present case is analogous to that of LTB collapse (the dependence of the visibility of the singularity on the initial data is exactly the same), we do not expect the inclusion of $k(r) \neq 0$ to qualitatively change the end result of collapse. Secondly, a more general issue—that concerns not only the present analysis, but the issue of generic gravitational collapse—is that of the physical reasonability of ‘dust’. Whilst early studies of high-density nuclear matter suggested that such an approximation could be a legitimate one

(at least for spherical collapse) [40], it is now well known that radial and tangential pressures *must* be taken into account at the very late stages of realistic astrophysical collapse [41]. The inclusion of radial pressure in spherical dust collapse has recently been considered [42], and it has been found to cover part (but not all) of the singularity spectrum, i.e. configurations that would otherwise develop locally naked singularities, end up in a black hole. The inclusion of tangential pressure has been considered for special cases of the Einstein cluster class [14,43] and for more generic configurations [44], with markedly different results: tangential stresses tend to uncover part of the singularity spectrum, i.e. configurations that would otherwise terminate in a black hole, develop a locally naked singularity. For the reasons mentioned above, we expect the inclusion of radial and tangential stresses in quasi-spherical asymptotically de Sitter dust collapse to produce similar effects to those of the spherical case.

Unlike the matter content, however, the effects of non-sphericity on the end state of collapse are much less clear, and further analyses, of stronger deviations from spherical symmetry, need to be undertaken to confidently establish the role of the latter in the final state of generic gravitational collapse.

Acknowledgments

It is a pleasure to thank J Barrow and K Thorne for helpful comments and discussions, and F Mena and B Nolan for privately communicating their recent results on geodesics in spherical collapse [45], which led to a correction in this manuscript. This work was supported by FCT (Portugal) grant PRAXIS XXI-BPD-16301-98, NSF grant AST-9731698 and by NSF grant PHY-0099568.

References

- [1] Gonçalves S M C V 2001 *Phys. Rev. D* **63** 064017
- [2] Deshingkar S S, Chamorro A, Jhingan S and Joshi P S 2001 *Phys. Rev. D* **63** 124005
- [3] Joshi P S and Singh T P 1993 *Phys. Rev. D* **47** 5357
- [4] Joshi P S and Dwivedi I H 1994 *Commun. Math. Phys.* **166** 117
- [5] Joshi P S and Dwivedi I H 1999 *Class. Quantum Grav.* **16** 41
- [6] Lake K 1992 *Phys. Rev. Lett.* **68** 3129
- [7] Ori A and Piran T 1987 *Phys. Rev. Lett.* **59** 2137
- [8] Harada T 1998 *Phys. Rev. D* **58** 104015
- [9] Thorne K S 1965 *Phys. Rev. B* **139** 251
- [10] Thorne K S 1972 *Magic Without Magic: John Archibald Wheeler* ed J Klauder (San Francisco, CA: Freeman)
- [11] Shapiro S L and Teukolsky S A 1991 *Phys. Rev. Lett.* **66** 994
- [12] Shapiro S L and Teukolsky S A 1992 *Phys. Rev. D* **45** 2006
- [13] Nakamura T, Shibata M and Nakao K 1993 *Prog. Theor. Phys.* **89** 821
- [14] Harada T, Iguchi H and Nakao K 1998 *Phys. Rev. D* **58** 041502
- [15] Iguchi H, Harada T and Nakao K 1999 *Prog. Theor. Phys.* **101** 1235
- [16] Iguchi H, Harada T and Nakao K 2000 *Prog. Theor. Phys.* **103** 53
- [17] Barve S, Singh T P, Vaz C and Witten L 1999 *Class. Quantum Grav.* **16** 1727
- [18] Szekeres P 1975 *Commun. Math. Phys.* **41** 55
- [19] Szekeres P 1975 *Phys. Rev. D* **12** 2941
- [20] Joshi P S and Królak A 1996 *Class. Quantum Grav.* **13** 3069
- [21] Deshingkar S S, Jhingan S and Joshi P S 1998 *Gen. Rel. Grav.* **30** 1477
- [22] Riess A G *et al* 1998 *Astron. J.* **116** 1009
- [23] Perlmutter S *et al* 1999 *Astrophys. J.* **517** 565
- [24] Zehani I and Dekel A 1999 *Nature* **401** 252
- [25] Markovic D and Shapiro S L 2000 *Phys. Rev. D* **61** 084029
- [26] Barrow J D and Stein-Schabes J 1984 *Phys. Lett. A* **103** 315
- [27] Bonnor W B 1976 *Commun. Math. Phys.* **51** 191

- [28] Yodzis P, Seifert H J and zum Hagen H M 1973 *Commun. Math. Phys.* **34** 135
- [29] Gonçalves S M C V 2001 *Phys. Rev. D* **63** 124017
- [30] Newman R P A C 1986 *Class. Quantum Grav.* **3** 527
- [31] Papapetrou A and Hamoui A 1967 *Ann. Inst. H Poincaré* **VI** 343
- [32] Abramovitz M and Stegun I A 1964 *Handbook of Mathematical Functions* (New York: Dover)
- [33] Hawking S W and Ellis G F R 1973 *The Large Scale Structure of Space-Time* (Cambridge: Cambridge University Press)
- [34] Christodoulou D 1984 *Commun. Math. Phys.* **93** 171
- [35] Kevorkian J 1999 *Partial Differential Equations: Analytical Solution Techniques* (New York: Springer)
- [36] Tipler F J 1977 *Phys. Lett. A* **64** 8
- [37] Clarke C J S and Królak A 1985 *J. Geom. Phys.* **2** 127
- [38] Ori A 2000 *Phys. Rev. D* **61** 064016
- [39] Jhingan S and Joshi P S 1997 The structure of singularity in spherical inhomogeneous dust collapse *Preprint gr-qc/9701016*
- [40] Hagedorn R 1965 *Nuovo Cimento Suppl.* **3** 147
- [41] Miller J C and Sciamia D W 1980 *General Relativity and Gravitation* ed A Held (New York: Plenum) vol 2
- [42] Gonçalves S M C V and Jhingan S 2001 Singularities in gravitational collapse with radial pressure *Gen. Rel. Grav.* at press
(Gonçalves S M C V and Jhingan S 2001 *Preprint gr-qc/0107054*)
- [43] Jhingan S and Magli G 2000 *Phys. Rev. D* **61** 124006
- [44] Gonçalves S M C V, Jhingan S and Magli G 2001 The spectrum of endstates of gravitational collapse with tangential stresses *Phys. Rev. D* submitted
- [45] Mena F C and Nolan B C 2001 Non-radial null geodesics in spherical dust collapse *Class. Quantum Grav.* **18** 4531
(Mena F C and Nolan B C 2001 *Preprint gr-qc/0108008*)